

Surgical Rehabilitation Protocol Patellofemoral Arthroplasty

The following information is to help make your recovery from surgery as smooth and rapid as possible. If you have any questions or concerns, contact Dr. Mayo's team at the number above. You will have appointments with Dr. Mayo at ~1-2 weeks postop.

Phase 1: Recovery from Surgery – 0-4 Weeks After Surgery

Goals	<ul style="list-style-type: none"> • Educate patient on rehab progression • Diminish joint swelling and pain • Restore patellar mobility • Restore full passive knee extension • Gradually improve knee flexion to 90 degrees • Re-establish quadriceps control and activation • Restore independent ambulation within precautions
Precautions	<ul style="list-style-type: none"> • Brace: Bledsoe on when ambulating and sleeping. • Weight Bearing: Weight bearing as tolerated in brace with crutches, wean from crutches when gaining quad control • Range of Motion: 0-120 degrees. Do not put a pillow until knee for comfort. Place pillow under ankle to keep knee in terminal extension. • Wound Care: No swimming or submerging in water until wounds healed • If performed in conjunction with meniscal repair or other surgery, defer to most restrictive guidelines • Call Dr. Mayo if: Significant wound drainage or dehiscence, purulence, erythema.
Therapeutic Exercises <i>See last page for example exercises</i>	<ul style="list-style-type: none"> • Strengthening: Quad sets, four-way SLR, ankle pumps, closed chain (mini squats, heel raise, etc.) • Proprioception: Weight shifting • Conditioning: Stationary bike, no resistance • Modalities: BFR, NMES • Manual Therapy: Patella and soft tissue mobilization, passive knee flexion to 120 degrees
Home Instructions	<ul style="list-style-type: none"> • Wound Care: Remove large bulky dressing on postoperative day 3. Leave large bandage in place. Sutures will be removed at ~2 weeks in clinic. • Bathing: Showering permitted once after bulky dressing removed. No submerging in water (bath/pool/lake/etc.) for 4 weeks. • Driving: Must be off all narcotic pain meds when operating vehicle <ul style="list-style-type: none"> ○ 1 week for automatic cars, left leg surgery ○ 2-4 weeks for standard/manual cars or right leg surgery • Sleeping: Sleep with brace locked in extension for 1 week • Ice and Elevation: Ice for 20 minutes every hour for the first week, elevate leg with knee in full extension as much as possible. Ice as needed after 1 week. • Home Exercise: As instructed by physical therapy.
Criteria to Progress	<ul style="list-style-type: none"> <input type="checkbox"/> Knee ROM: 0-90 degrees <input type="checkbox"/> Perform SLR without quad lag and good quad activation <input type="checkbox"/> Normalized gait per precautions <input type="checkbox"/> Normal patellar mobility <input type="checkbox"/> Minimal swelling/inflammation



Surgical Rehabilitation Protocol Patellofemoral Arthroplasty

The following information is to help make your recovery from surgery as smooth and rapid as possible. If you have any questions or concerns, contact Dr. Mayo's team at the number above. You will have appointments with Dr. Mayo at ~6 weeks postop

Phase 2: Early Strengthening/Neuromuscular Control – 4-8 Weeks After Surgery

Goals	<ul style="list-style-type: none"> • Eliminate inflammation and swelling • Full knee ROM (0-135 degrees) or hyperextension if normal • Normal gait on all surfaces without brace or assistive device • Improve lower extremity strength, proprioception, balance, neuromuscular control, and confidence • Demonstrate stability with dynamic knee activities (no varus/valgus deviations)
Precautions	<ul style="list-style-type: none"> • Brace: Transition out of brace as gait is normalized • Weight Bearing: Full weight bearing • Range of Motion: No restrictions • Call Dr. Mayo if: Not achieving full range of motion, large effusion
Therapeutic Exercises <i>See last page for example exercises</i>	<ul style="list-style-type: none"> • Strengthening: Advancement of multi-plane closed chain activities, open chain, core strengthening and functional standing activities • Proprioception: One leg balance, wobble boards, BAPS • Conditioning: Walking program • Modalities: BFR, NMES • Manual Therapy: Patella and soft tissue mobilization, passive knee flexion to 135 degrees, prone quadriceps stretching, joint mobilization as needed
Home Instructions	<ul style="list-style-type: none"> • Driving: OK to drive assuming off narcotic pain medication • Sleeping: OK to remove brace • Ice and Elevation: Ice as needed for pain and swelling after activity • Home Exercises: As instructed by physical therapy
Criteria to Progress	<ul style="list-style-type: none"> <input type="checkbox"/> Full knee ROM <input type="checkbox"/> Minimal swelling/inflammation <input type="checkbox"/> No pain with exercises <input type="checkbox"/> Normal gait on all surfaces at community level distances



Surgical Rehabilitation Protocol Patellofemoral Arthroplasty

The following information is to help make your recovery from surgery as smooth and rapid as possible. If you have any questions or concerns, contact Dr. Mayo's team at the number above. You will have appointments with Dr. Mayo at ~3 months postop.

Phase 3: Advanced Activity Phase – 8-12 Weeks After Surgery

Goals	<ul style="list-style-type: none"> • Normalize lower extremity strength to >85% non-involved extremity • Enhance muscular power and endurance • Improve neuromuscular control and proprioception exercises • Perform selected sport-specific drills • Initiate plyometric exercises • Improve aerobic endurance • Physician clearance to initiate return to running and functional progression
Precautions	<ul style="list-style-type: none"> • Brace: None • Call Dr. Mayo if: Increased effusion, not full range of motion
Therapeutic Exercises	<ul style="list-style-type: none"> • Phase 2 exercises • Elliptical and other low impact aerobic activities <p><i>See last page for example exercises</i></p>
Home Instructions	<ul style="list-style-type: none"> • Home exercises: Workouts in gym, focus per physical therapist
Criteria to Progress	<ul style="list-style-type: none"> <input type="checkbox"/> No pain with activities of daily living or work specific movement <input type="checkbox"/> Good knee control with single leg dynamic proprioceptive activities



Surgical Rehabilitation Protocol Patellofemoral Arthroplasty

The following information is to help make your recovery from surgery as smooth and rapid as possible. If you have any questions or concerns, contact Dr. Mayo's team at the number above. You will have appointments with Dr. Mayo at ~6 months and 1 year postop.

Phase 4: Return to Activity – 12+ Weeks after Surgery to Return

Goals	<ul style="list-style-type: none"> • Improve muscular strength and endurance • Good control and no pain with all activities of daily living as well as work specific movements • Avoid deep squatting, lunging or kneeling • Return to ADLs • Progress proprioception
Precautions	<ul style="list-style-type: none"> • Brace: No brace recommended • No return to full participation in recreational activities or work until cleared by Dr. Mayo
Therapeutic Exercises	<ul style="list-style-type: none"> • Continue all exercises • Plyometric drills • Initiate running • Advanced proprioception drills
Home Instructions	<ul style="list-style-type: none"> • Home Exercises: Workouts in gym, focus per physical therapist
Criteria for Full Return to Activity	<input type="checkbox"/> Able to perform all required activities safely and without pain



Surgical Rehabilitation Protocol Patellofemoral Arthroplasty

Sample Rehabilitation Exercises by Phase

Phase I	Phase II
Week 0-4	Week 4-12
<p style="text-align: center;"><u>Week 0-2:</u></p> <p>Range of Motion</p> <ul style="list-style-type: none"> • Ankle pumps • Overpressure into full, passive knee extension • Active and passive gradual knee flexion 0-90 degree • Hamstring and gastrocnemius stretch • Heel slides, AAROM prone knee flexion, seated flexion stretch <p>Strengthening</p> <ul style="list-style-type: none"> • Straight leg raises (Flexion, Abduction, Adduction) • Quadriceps activation sets and stimulation • Calf strengthening • Closed kinetic chain exercises: mini squat and weight shifts Closed kinetic chain quad strengthening (wall sits, step-ups, mini-squats, leg press) • Multi-angle isometric at 90-60 degree extension • Knee extension 90-40 degrees <p>Manual Therapy/Modalities</p> <ul style="list-style-type: none"> • NMES is strongly recommended • Stationary bike for ROM/strength no resistance • Patellar mobilization • Blood flow restriction <p style="text-align: center;"><u>Week 2-4</u></p> <ul style="list-style-type: none"> • Continue exercises from week 1 and 2 <p>Range of Motion</p> <ul style="list-style-type: none"> • Passive range of motion from 0-100 <p>Strengthening</p> <ul style="list-style-type: none"> • Hip straight leg raises in 4 planes (in brace until can perform without quad lag) • Hamstring bridge • Leg press 0-60 degrees • Half squats 0-40 degrees • Front and side lunges • Progressive resistance extension program starting with 1lb progress 1lb a week • OKC passive/active joint repositioning 90,60, 30 degrees • CKC joint repositioning during squat/lunges <p>Proprioception/Plyometric</p> <ul style="list-style-type: none"> • Initiation squat on tilt board • Initiate proprioceptive exercises (single leg balance, ball toss, balance beam, BOSU, Airex) 	<p style="text-align: center;"><u>Week 4-7:</u></p> <p>Range of Motion</p> <ul style="list-style-type: none"> • Advance ROM <p>Strengthening</p> <ul style="list-style-type: none"> • Progress isometric strengthening program • Leg press (0-100 degrees) • Knee extension 90 to 40 degrees - Avoid open chain quad strengthening 30-0 degrees • Passive/active reposition OKC • Hamstring Curls (isotonics) • Hip Abduction and Adduction • Hip Flexion and Extension • Lateral Step-Overs, Lunges (straight plane and multi-plane drills) • Lateral Step Ups, Front Step Downs • Wall slides/squats • Vertical Squats • Standing or Seated Toe Calf Raises <p>Endurance</p> <ul style="list-style-type: none"> • Bicycle and Stair Stepper Machine • Pool Program (Running, agility, Backward Running, Hip and Leg Exercises) • Unloading treadmill walking <p>Proprioception/Plyometric</p> <ul style="list-style-type: none"> • Proprioception Drills • Biodex Stability System (Balance, Squats, etc) • Progress to balance and ball throws • Tilt board repositioning, balance, and squats (perturbation) <p style="text-align: center;"><u>Week 8-12</u></p> <ul style="list-style-type: none"> • Continue all exercises listed in Weeks 4-6 <p>Strengthening</p> <ul style="list-style-type: none"> • Leg Press Sets (single leg) 0-100 degrees and 40-100 degrees • Isokinetic exercises (90 to 40 degrees) (120 to 240 degrees/second) <p>Proprioception/Plyometric</p> <ul style="list-style-type: none"> • Plyometric Leg Press • Biodex stability system • Training on tilt board • Perturbation Training <p>Endurance</p> <ul style="list-style-type: none"> • Walking Program • Bicycle/Stair Stepper/Elliptical Machine for endurance • May initiate running program (weeks 10-12) (Physician Decision) with brace



Surgical Rehabilitation Protocol Patellofemoral Arthroplasty

Phase III	Phase IV
Weeks 12-24	Weeks 24+
<p><u>Strengthening</u></p> <ul style="list-style-type: none"> • Continue strengthening - advance resistance and repetitions (ball hamstring curls, single leg press, core stabilization) • Continue all strengthening drills • Leg press • Wall squats • Hip Abd/Adduction • Hip Flex/Ext • Knee Extension 90-40 • Hamstring curls • Standing toe calf raises • Step down • Lateral step ups • Lateral lunges • Plyometric leg press • Neuromuscular training • Lateral step-overs cones • Tilt board drills • May initiate lateral agility drills • Backward running • Spin bike • Cybex training <p><u>Proprioception/Plyometric</u></p> <ul style="list-style-type: none"> • Pre-running exercises (low skips, punch steps, double punch steps, hurdle walks, high skips, kickbacks, step-overs) • Advance proprioceptive exercises (BOSU, single leg dynamic balance, dual task balance) • Agility drills (ladder, side shuffles, crossovers, backwards run, quick start/stops, zig-zags, cutting) • Jump training (shuttle training, trampoline, landing technique, box jumps, single leg hops, tuck jumps) • Return to running – treadmill, with transition to level outdoor surfaces <p><u>Endurance/Sport Specific</u></p> <ul style="list-style-type: none"> • Initiate running program (weeks 10-12) (Physician Decision) with brace • May initiate light sport program (golf) (Surgeon Decision) with brace • 	<ul style="list-style-type: none"> • Continue strengthening exercises • Continue neuromuscular control drills • Continue plyometrics drills • Progress running and agility program • Gradually progress level of participation in sport specific training • Running/cutting/agility drills • Gradual return to sport drills • Running on all surfaces

